


## 6.3 Polar Coordinates

In this section we will learn a new coordinate system. In this system we plot a point in the form  $(r, \theta)$ . As shown in the picture below you first draw angle  $\theta$  in standard form. Then you label how long  $r$  is:



EXAMPLE: Plot  $\left(4, \frac{3\pi}{4}\right)$  in the polar coordinate system.

EXAMPLE: Plot  $\left(5, \frac{5\pi}{3}\right)$  in the polar coordinate system.

EXAMPLE: Plot  $(-3, 120^\circ)$  in the polar coordinate system.

EXAMPLE: Plot  $\left(-3, -\frac{\pi}{2}\right)$  in the polar coordinate system.

EXAMPLE: Plot  $\left(-3, -\frac{3\pi}{4}\right)$  in the polar coordinate system.

### Equivalent Polar Coordinates

There are more than one way to arrive at the same angle. For example in the previous problem,  $-135$  degrees is the same as  $360^\circ + (-135^\circ) = 225^\circ$ . If we have  $120$  degrees then this is the same as  $120^\circ + (-360^\circ) = -240^\circ$ . So for negative angles, just add  $360$  degrees. For positive angles add negative  $360$  degrees to find the equivalent angle. So basically we can either move clockwise or counterclockwise to arrive at the same angle.

$$\begin{aligned}(r, \theta) &= (r, \theta \pm 2\pi) \text{ or } (r, \theta) = (r, \theta \pm 360^\circ) \\ (r, \theta) &= (-r, \theta \pm \pi) \text{ or } (r, \theta) = (-r, \theta \pm 180^\circ)\end{aligned}$$

EXAMPLE: Given the polar coordinate  $(5, 300^\circ)$ , find an equivalent polar coordinate that has the following characteristics: a.)  $-360^\circ \leq \theta \leq 0$ ,  $r > 0$  b.)  $0 \leq \theta \leq 360^\circ$ ,  $r < 0$ , c.)  $360^\circ \leq \theta \leq 720^\circ$ ,  $r > 0$

a.)  $-360^\circ \leq \theta \leq 0$ ,  $r > 0$

$$\text{b.) } 0 \leq \theta \leq 360^\circ, r < 0$$

$$\text{c.) } 360^\circ \leq \theta \leq 720^\circ, r > 0$$

EXAMPLE: Given the polar coordinate  $\left(4, \frac{3\pi}{4}\right)$ , find an equivalent polar coordinate that has the following characteristics: a.)  $-2\pi \leq \theta \leq 0, r > 0$  b.)  $0 \leq \theta \leq 2\pi, r < 0$ , c.)  $2\pi \leq \theta \leq 4\pi, r > 0$

$$\text{a.) } -2\pi \leq \theta \leq 0, r > 0$$

$$\text{b.) } 0 \leq \theta \leq 2\pi, r < 0$$

$$\text{c.) } 2\pi \leq \theta \leq 4\pi, r > 0$$

EXAMPLE: Given the polar coordinate  $(-2, -120^\circ)$ , find an equivalent polar coordinate that has the following characteristics: a.)  $-360^\circ \leq \theta \leq 0$ ,  $r > 0$  b.)  $0 \leq \theta \leq 360^\circ$ ,  $r < 0$ , c.)  $360^\circ \leq \theta \leq 720^\circ$ ,  $r > 0$

a.)  $-360^\circ \leq \theta \leq 0$ ,  $r > 0$

b.)  $0 \leq \theta \leq 360^\circ$ ,  $r < 0$

c.)  $360^\circ \leq \theta \leq 720^\circ$ ,  $r > 0$

**Conversion formulas from polar to rectangular coordinates.**

$$x = r \cos \theta$$

$$y = r \sin \theta$$

$$x^2 + y^2 = r^2$$

EXAMPLE: Convert  $\left(5, \frac{\pi}{3}\right)$  into a rectangular point.

EXAMPLE: Convert  $\left(-3, -\frac{\pi}{4}\right)$  into a rectangular point.

EXAMPLE: Convert  $\left(-2, \frac{2\pi}{3}\right)$  into a rectangular point.

EXAMPLE: Convert the equation  $r = 5 \sec \theta$  into a rectangular equation.

EXAMPLE: Convert the equation  $r = 4$  into a rectangular equation.

EXAMPLE: Convert the equation  $r = 2 \sin \theta - 4 \cos \theta$  into a rectangular equation.

### Conversion formulas from rectangular to polar coordinates

$$x^2 + y^2 = r^2$$

If  $(x, y)$  is in the first or fourth quadrant, then  $\theta = \tan^{-1} \frac{y}{x}$ .

If  $(x, y)$  is in the second or third quadrant, then  $\theta = \tan^{-1} \frac{y}{x} + \pi$ .

EXAMPLE: Convert  $(-3, 3)$  into a polar coordinate. Express your angle in radians.

EXAMPLE: Convert  $(-2, -2\sqrt{3})$  into a polar coordinate. Express your angle in radians.

EXAMPLE: Convert  $\left(\frac{1}{2}, -\frac{\sqrt{3}}{2}\right)$  into a polar coordinate. Express your angle in radians.

EXAMPLE: Convert the equation  $x^2 + y^2 = x$  into a polar equation and solve for  $r$ .

EXAMPLE: Convert the equation  $4x^2y = 1$  into a polar equation and solve for  $r$ .

EXAMPLE: Convert the equation  $y^2 = 2x$  into a polar equation and solve for  $r$ .