

2.4 Continuity



Continuity: graph is connected with no breaks or holes.

Let $f(x)$ be a graph. Continuity at $x = c$ occurs if all three of the following are true:


- 1.) $f(c)$ is defined. (There are no vertical asymptotes or holes at $x = c$)
- 2.) $\lim_{x \rightarrow c} f(x)$ exists. (If it didn't exist, then there must be a break in the graph or a vertical asymptote.)
- 3.) $\lim_{x \rightarrow c} f(x) = f(c)$ (Since these y-values are equal, there can't be a hole.)

If even one of the above is false then the function is discontinuous. There are two types of discontinuities:

- 1.) Removable discontinuity: A point can be assigned to "plug up" the hole.
- 2.) Non-removable discontinuity: No point can cause the graph to be connected.



Let's look at the below graph from the previous section. Can you tell which places the graph is discontinuous? Which ones are removable and non-removable?



We see that the graph will be discontinuous at $x = -2, 0, 2,$ and 4 . The only removable discontinuity is at $x = 2$. The others are non-removable.

EXAMPLE: Describe the set of x -values where the function is continuous, using interval notation.

$$f(x) = (3 - x)^{\frac{1}{5}}$$

For all these problems you want to look for any place where the equation will be undefined. This only occurs if you are dividing by a negative number or taking the even root of a negative number. In this problem there are no places where we are dividing by zero, and we have an odd root here since $f(x)$ can be written as:

$f(x) = \sqrt[5]{3 - x}$. Therefore, it will be continuous at every value of x . So our answer is $(-\infty, \infty)$.

EXAMPLE: Describe the set of x-values where the function is continuous, using interval notation.

$$f(x) = \sqrt{6x - 35}$$

We are only allowed to take the square root of positive numbers or zero. Therefore this is the only place where the graph is continuous. So to solve this one, we will let everything inside the square root be greater than or

equal to zero: $6x - 35 \geq 0$. Solving this will give us $x \geq \frac{35}{6}$. In interval notation the answer is: $\left[\frac{35}{6}, \infty\right)$.

EXAMPLE: Describe the set of x-values where the function is continuous, using interval notation.

$$f(x) = \frac{1}{x^2 + 1}$$

In the problem above, since no x value will make the bottom zero then this graph will be continuous for all x, therefore the answer is $(-\infty, \infty)$.

EXAMPLE: Determine the x-values where $f(x) = \frac{\tan x}{x^2 + 1}$ is continuous. Write an exact answer, using π as needed. Write the expression using n , where n is any **odd** integer.

The function $f(x) = \frac{\tan x}{x^2 + 1}$ is continuous on $(-\infty, \infty)$ except at $x = \underline{\hspace{2cm}}$

From the previous example we know that the bottom won't ever be zero. However, we still need to consider the top. We need to find where tangent could be undefined. Identities tell us that $\tan x = \frac{\sin x}{\cos x}$ so we need to see

where $\cos x = 0$. This happens at $x = \frac{\pi}{2}, \frac{3\pi}{2}, \frac{5\pi}{2}, \dots$. We can generalize this by stating that $f(x)$ will be

discontinuous at $x = \frac{n\pi}{2}$, where n is odd. So the function is continuous on $(-\infty, \infty)$ except at $x = \frac{n\pi}{2}$.

EXAMPLE: Determine the x-values where $f(x) = 4 \csc(4x)$ is continuous. Write an exact answer, using π as needed. Write the expression using n , where n is any integer.

The function $f(x) = 4 \csc(4x)$ is continuous on $(-\infty, \infty)$ except at $x = \underline{\hspace{2cm}}$

This one can be written with identities: $f(x) = \frac{4}{\sin(4x)}$. We need to see where $\sin(4x) = 0$. Suppose the $4x$

was just x . Then we know that would be zero at $0, \pi, 2\pi, \dots$. However, we have $4x$ so basically

$4x = 0, \pi, 2\pi, 3\pi, 4\pi, \dots$. Therefore by solving for x we get $x = 0, \frac{\pi}{4}, \frac{\pi}{2}, \frac{3\pi}{4}, \pi, \dots$. So this is discontinuous at

$x = \frac{n\pi}{4}$ where n is any integer. These would be considered non-removable discontinuities since these are

vertical asymptotes. So the function is continuous on $(-\infty, \infty)$ except at $x = \frac{n\pi}{4}$.

EXAMPLE: Indicate the x -values (if any) at which f is not continuous. Then indicate the x -values where f has removable discontinuities and nonremovable discontinuities.

$$f(x) = \frac{x-3}{x^2-9}$$

Before doing anything we want to factor: $f(x) = \frac{x-3}{(x+3)(x-3)}$. We see that 3 and -3 will make the bottom zero, so these are the discontinuities. Let's classify each discontinuity. If you can cancel a term from the top and bottom we will have $f(x) = \frac{1}{x+3}$ in which there will be a vertical asymptote at $x = -3$. This means that there is a nonremovable discontinuity at $x = -3$. What about 3? This is actually a hole in the graph so $x = 3$ is a removable discontinuity.

EXAMPLE: Indicate the x -values (if any) at which f is not continuous. Then indicate the x -values where f has removable discontinuities and nonremovable discontinuities.

$$f(x) = \frac{x+2}{x^2-2x-15}$$

Like in the previous example we first want to factor the top and bottom.

We will get: $f(x) = \frac{x+2}{(x+3)(x-5)}$. We see that $x = -3$ and $x = 5$

would be all the discontinuities. This time nothing cancels, so both

$x = -3$ and $x = 5$ would be considered nonremovable discontinuities.

